

Integrable Rosochatius deformations of higher-order constrained flows and the soliton hierarchy with self-consistent sources

Yubin Yao¹ and Yunbo Zeng²

Department of Mathematics, Tsinghua University, Beijing 100084, PR China

Abstract We propose a systematic method to generalize the integrable Rosochatius deformations for finite dimensional integrable Hamiltonian systems to integrable Rosochatius deformations for infinite dimensional integrable equations. Infinite number of the integrable Rosochatius deformed higher-order constrained flows of some soliton hierarchies, which includes the generalized integrable Hénon-Heiles system, and the integrable Rosochatius deformations of the KdV hierarchy with self-consistent sources, of the AKNS hierarchy with self-consistent sources and of the mKdV hierarchy with self-consistent sources as well as their Lax representations are presented.

1 Introduction

Rosochatius found that it would still keep the integrability to add a potential of the sum of inverse squares of the coordinates to that of the Neumann system [1, 2]. The deformed system is called Neumann-Rosochatius system. Wojciechowski obtained an analogy system for the Garnier system as a stationary KdV flow in 1985 [3, 4]. In 1997, Kubo et al. [5] constructed the analogy system for the Jacobi system [6] and the geodesic flow equation on the ellipsoid based upon the Deift technique and a theorem that the Gauss map transforms the Neumann system to the Jacobi system [7, 8]. All of these systems have the same character that they are integrable Hamiltonian systems containing N arbitrary parameters and the original finite dimensional integrable Hamiltonian systems are recovered when all these parameters vanish. In fact, these systems are a sort of integrable deformations of the corresponding integrable Hamiltonian systems, which are called integrable Rosochatius deformation. The resulting systems are called the Rosochatius-type integrable systems. These systems have important physical applications. For examples, Neumann-Rosochatius system can be used to describe the dynamics of rotating closed string solutions in $AdS_5 \times S^5$ and the membranes on $AdS_4 \times S^7$ [9]- [17]. The Garnier-Rosochatius system can be used to solve the multicomponent coupled nonlinear Schrödinger equation [3, 4, 18]. It is not difficult to see that each of the above system has its own origin. In Ref. [19], Zhou generalize the Rosochatius method to study the integrable Rosochatius deformations of some explicit constrained flows of soliton equations.

However, so far the Rosochatius deformations are limited to few finite dimensional integrable Hamiltonian systems(FDIHSs). It is natural to ask whether there exist integrable Rosochatius deformations for infinite dimensional integrable equations. The main

¹yqyao@math.tsinghua.edu.cn

²yzeng@math.tsinghua.edu.cn

purpose of this paper is to generalize the Rosochatius deformation from FDIHSs to infinite dimensional integrable equations. We will investigate the integrable Rosochatius deformations firstly for infinite number of higher-order constrained flows of some soliton equations, then for some soliton hierarchies with self-consistent sources.

In recent years the constrained flows of soliton equations obtained from the symmetry reduction of soliton equations, which can be transformed into a FDIHS, attracted a lot of attention [20]- [33]. Many well-known FDIHSs are recovered by means of constrained flows. Furthermore, soliton equation can be factorized into two commuting constrained flows, which provides an effective way to solve the soliton equations through solving the constrained flows.

To make the paper self-contained, we first briefly recall the higher-order constrained flows of the soliton hierarchy and the soliton hierarchy with self-consistent sources. Consider a hierarchy of soliton equation which can be formulated as an infinite dimensional Hamiltonian systems

$$u_{t_n} = J \frac{\delta H_n}{\delta u}. \quad (1)$$

The auxiliary linear problems associated with (1) are given by

$$\phi_x = U(\lambda, u)\phi, \quad \phi = (\phi_1, \phi_2)^T \quad (2a)$$

$$\phi_{t_n} = V^{(n)}(\lambda, u)\phi. \quad (2b)$$

Then the higher-order constrained flows of (1) consist of the equations obtained from the spectral problem (2a) for N distinct λ_j and the restriction of the variational derivatives for the conserved quantities H_n and λ_j [23]

$$J\left[\frac{\delta H_n}{\delta u} + \sum_{j=1}^N \frac{\delta \lambda_j}{\delta u}\right] = 0, \quad (3a)$$

$$\phi_{j,x} = U(\lambda_j, u)\phi_j, \quad \phi_j = (\phi_{1j}, \phi_{2j})^T, \quad j = 1, 2, \dots, N. \quad (3b)$$

By introduction the so called Jacobi-Ostrogradsky coordinates [34], (3) can be transformed into a FDIHS. The Lax representation of (3) can be deduced from the adjoint representation of (2) [24]

$$N_x^{(n)} = [U, N^{(n)}]. \quad (4)$$

Mainly, $N^{(n)}$ has the following forms

$$N^{(n)}(\lambda) = V^{(n)}(\lambda) + N_0 = \begin{pmatrix} A(\lambda) & B(\lambda) \\ C(\lambda) & -A(\lambda) \end{pmatrix},$$

where

$$N_0 = \sum_{j=1}^N \frac{1}{\lambda - \lambda_j} \begin{pmatrix} \phi_{1j}\phi_{2j} & -\phi_{1j}^2 \\ \phi_{2j}^2 & -\phi_{1j}\phi_{2j} \end{pmatrix}, \quad or \quad N_0 = \sum_{j=1}^N \frac{1}{\lambda^2 - \lambda_j^2} \begin{pmatrix} \lambda\phi_{1j}\phi_{2j} & -\lambda_j\phi_{1j}^2 \\ \lambda_j\phi_{2j}^2 & -\lambda\phi_{1j}\phi_{2j} \end{pmatrix}.$$

The soliton hierarchy with self-consistent sources is defined by [25, 26]

$$u_{t_n} = J \left[\frac{\delta H_n}{\delta u} + \sum_{j=1}^N \frac{\delta \lambda_j}{\delta u} \right], \quad (5a)$$

$$\phi_{j,x} = U(\lambda_j, u) \phi_j, \quad j = 1, 2, \dots, N. \quad (5b)$$

Since the higher-order constrained flows (3) are just the stationary equations of (5), the zero-curvature representation for (5) can be induced from (4) as follows

$$U_{t_n} - N_x^{(n)} + [U, N^{(n)}] = 0 \quad (6)$$

which implies that (5) is Lax integrable. In fact, (5) can also be formulated as an infinite-dimensional integrable Hamiltonian system with t-type Hamiltonian operator by taking t as the 'spatial' variable and x as the evolution parameter as well as introducing the Jacobi-Ostrogradsky coordinates [35, 36]. Then we can find the Poisson bracket defined by the t-type Hamiltonian operator, and the conserved density when considering x as the evolution parameter. The t-type bi-Hamiltonian description for KdV equation with self-consistent sources and for Jaulent-Miodek equation with self-consistent sources were presented in [35] and [36], respectively.

The soliton equation with self-consistent sources have important physical application, for example, the KdV equation with self-consistent sources describes the interaction of long and short capillary-gravity wave [37]- [39].

In this paper, in the same way [19], we construct the Rosochatius deformation $\tilde{N}^{(n)}$ of Lax matrix $N^{(n)}$ by replacing ϕ_{2j}^2 in the matrix N_0 with $\phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2}$, namely the entries of $\tilde{N}^{(n)}$ are given by

$$\tilde{A}(\lambda) = A(\lambda), \quad \tilde{B}(\lambda) = B(\lambda), \quad \tilde{C}(\lambda) = C(\lambda) + \sum_{j=1}^N \frac{\mu_j}{(\lambda - \lambda_j) \phi_{1j}^2},$$

or

$$\tilde{C}(\lambda) = C(\lambda) + \sum_{j=1}^N \frac{\lambda_j \mu_j}{(\lambda^2 - \lambda_j^2) \phi_{1j}^2}, \quad (7)$$

Then Lax representation (4) with $N^{(n)}$ replaced by $\tilde{N}^{(n)}$ gives rise to the Rosochatius deformations of (3). The fact that such a substitute keeps the relations of the Poisson brackets of $A(\lambda)$, $B(\lambda)$ and $C(\lambda)$ guarantees the integrability of Rosochatius deformations of (3). In this way we can obtain infinite number of integrable Rosochatius deformed FDIHSs constructed from the Rosochatius deformed higher-order constrained flows of KdV hierarchy, AKNS hierarchy and mKdV hierarchy, respectively. Among these Rosochatius deformed FDIHSs, it needs to point out that the Rosochatius deformation of the first higher-order constrained flow of KdV hierarchy contains the well-known generalized integrable Hénon-Heiles system, and can be regarded as the integrable multidimensional extension of Hénon-Heiles system. Then, based on the Rosochatius deformed higher-order constrained flows, the Rosochatius deformations of the soliton hierarchy with self-consistent sources (RDSH-SCS) can be constructed through (6) with $N^{(n)}$ replaced by $\tilde{N}^{(n)}$. The integrability of the

RDSHSCS can be explained by the fact that the RDSHSCS possesses the zero-curvature representation (6) with $N^{(n)}$ replaced by $\tilde{N}^{(n)}$ and its stationary reduction is integrable Rosochatius deformations of (3). In these way, we construct Rosochatius deformations of KdV hierarchy with self-consistent sources (RDKdVHSCS), of AKNS hierarchy with self-consistent sources (RDAKNSHSCS) and of mKdV hierarchy with self-consistent sources (RDmKdVHSCS), as well as their zero-curvature representations. We notice that there are two kinds of RDSHSCS. For RDKdVHSCS and RDmKdVHSCS, the Rosochatius formed terms only appear in (5b). However for RDAKNSHSCS Rosochatius deformed terms occur in both (5a) and (5b).

The structure of the paper is as follows. In Sec.2, we present the Rosochatius deformation of the higher-order constrained flows of KdV hierarchy and RDKdVHSCS as well as their Lax representations. In Sec.3, we obtain the Rosochatius deformation of higher-order constrained flows of AKNS hierarchy and RDAKNSHSCS as well as their Lax representations. In Sec.4, we obtain the Rosochatius deformation of higher-order constrained flows of mKdV hierarchy and RDmKdVHSCS and their Lax representations. In Sec.5, a conclusion is made.

2 The Rosochatius deformed KdV hierarchy with self-consistent sources

Consider the Schrödinger equation [40]

$$\phi_{1xx} + (\lambda + u)\phi_1 = 0, \quad (8)$$

which can be written in the matrix form

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}_x = U \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}, \quad U = \begin{pmatrix} 0 & 1 \\ -\lambda - u & 0 \end{pmatrix}. \quad (9)$$

The adjoint representation of (9) reads

$$V_x = [U, V]. \quad (10)$$

Set

$$V = \sum_{i=1}^{\infty} \begin{pmatrix} a_i & b_i \\ c_i & -a_i \end{pmatrix} \lambda^{-i}. \quad (11)$$

Solving (10) yields

$$\begin{aligned} a_k &= -\frac{1}{2}b_{k,x}, \quad b_{k+1} = Lb_k = -\frac{1}{2}L^{k-1}u, \quad c_k = -\frac{1}{2}b_{k,xx} - b_{k+1} - b_k u, \\ a_0 &= b_0 = 0, \quad c_0 = -1, \quad a_1 = 0, \quad b_1 = 1, \quad c_1 = -\frac{1}{2}u, \quad a_2 = \frac{1}{4}u_x, \\ b_2 &= -\frac{1}{2}u, \quad c_2 = \frac{1}{8}(u_{xx} + u^2), \quad b_3 = \frac{1}{8}(u_{xx} + 3u^2), \dots \end{aligned} \quad (12)$$

where $L = -\frac{1}{4}\partial^2 - u + \frac{1}{2}\partial^{-1}u_x$, $\partial = \frac{\partial}{\partial x}$.

Set

$$V^{(n)} = \sum_{i=1}^n \begin{pmatrix} a_i & b_i \\ c_i & -a_i \end{pmatrix} \lambda^{n-i} + \begin{pmatrix} 0 & 0 \\ b_{n+1} & 0 \end{pmatrix}, \quad (13)$$

and take

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}_{t_n} = V^{(n)}(u, \lambda) \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}. \quad (14)$$

Then the compatibility of the Eqs.(9) and (14) gives rise to the KdV hierarchy

$$u_{t_n} = -2b_{n+1,x} \equiv \partial \frac{\delta H_n}{\delta u}, \quad n = 0, 1, \dots, \quad (15)$$

where $H_n = 4b_{n+2}/2n + 1$. We have

$$\frac{\delta \lambda}{\delta u} = \phi_1^2, \quad L\phi_1^2 = \lambda\phi_1^2. \quad (16)$$

The higher-order constrained flows of the KdV hierarchy is given by [23],

$$\frac{\delta H_n}{\delta u} - \alpha \sum_{j=1}^N \frac{\delta \lambda_j}{\delta u} \equiv -2b_{n+1} - \alpha \sum_{j=1}^N \phi_{1j}^2 = 0, \quad (17a)$$

$$\phi_{1j,x} = \phi_{2j}, \quad \phi_{2j,x} = -(\lambda_j + u)\phi_{1j}, \quad j = 1, 2, \dots, N. \quad (17b)$$

According to (12), (16) and (17), we find the Lax representation (4) for (17) with

$$N^{(n)} = \sum_{k=0}^n \begin{pmatrix} a_k & b_k \\ c_k & -a_k \end{pmatrix} \lambda^{n-k} + \frac{\alpha}{2} \sum_{j=1}^N \frac{1}{\lambda - \lambda_j} \begin{pmatrix} \phi_{1j}\phi_{2j} & -\phi_{1j}^2 \\ \phi_{2j}^2 & -\phi_{1j}\phi_{2j} \end{pmatrix}. \quad (18)$$

By taking the so-called Jacobi-Ostrogradsky coordinates [34]

$$q_i = u^{(i-1)}, \quad i = 1, \dots, n-1,$$

$$p_i = \frac{\delta H_n}{\delta u^{(i)}} = \sum_{l \geq 0} (-\partial)^l \frac{\partial H_n}{\partial u^{(i+l)}}$$

and setting

$$\Phi_1 = (\phi_{11}, \phi_{12}, \dots, \phi_{1N})^T, \quad \Phi_2 = (\phi_{21}, \phi_{22}, \dots, \phi_{2N})^T, \quad Q = (\phi_{11}, \phi_{12}, \dots, \phi_{1N}, q_1, \dots, q_{n-1})^T,$$

$$P = (\phi_{21}, \phi_{22}, \dots, \phi_{2N}, p_1, \dots, p_{n-1})^T, \quad \Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_N).$$

Eq.(17) with $\alpha = \frac{2}{4^n}$ can be transformed into a FDIHS [23]

$$Q_x = \frac{\partial H}{\partial P}, \quad P_x = -\frac{\partial H}{\partial Q}, \quad (19)$$

with

$$H = \sum_{i=1}^{n-1} q_{i,x} p_i - H_n + \frac{1}{2} \langle \Phi_2, \Phi_2 \rangle + \frac{1}{2} \langle \Lambda \Phi_1, \Phi_1 \rangle + \frac{1}{2} q_1 \langle \Phi_1, \Phi_1 \rangle,$$

where $\langle \rangle$ denotes the inner production in R^N . For example, (17) for $n = 0$, $\alpha = -2$ gives rise to the Neumann system [20], (17) for $n = 1$, $\alpha = 1$ leads to the Garnier system [3, 20]. When $n = 2$, the Eq.(17) for $\alpha = \frac{1}{8}$ gives the first higher-order constrained flow [23]

$$u_{xx} + 3u^2 = -\frac{1}{2} \sum_{j=1}^N \phi_{1j}^2 = -\frac{1}{2} \langle \Phi_1, \Phi_1 \rangle, \quad (20a)$$

$$\phi_{1j,x} = \phi_{2j}, \quad \phi_{2j,x} = -(\lambda_j + u)\phi_{1j}, \quad j = 1, 2, \dots, N. \quad (20b)$$

Let $q_1 = u$, $p_1 = u_x$, (20) becomes a FDIHS (19) with

$$H = \frac{1}{2} \langle \Phi_2, \Phi_2 \rangle + \frac{1}{2} \langle \Lambda \Phi_1, \Phi_1 \rangle + \frac{1}{2} q_1 \langle \Phi_1, \Phi_1 \rangle + \frac{1}{2} p_1^2 + q_1^3,$$

and has the Lax representation (4) with the entries of $N^{(2)}$ given by

$$A(\lambda) = \frac{1}{4} p_1 + \frac{1}{16} \sum_{j=1}^N \frac{\phi_{1j} \phi_{2j}}{\lambda - \lambda_j}, \quad B(\lambda) = \lambda - \frac{1}{2} q_1 - \frac{1}{16} \sum_{j=1}^N \frac{\phi_{1j}^2}{\lambda - \lambda_j},$$

$$C(\lambda) = \lambda^2 - \frac{q_1}{2} \lambda - \frac{q_1^2}{4} - \frac{1}{16} \langle \Phi_1, \Phi_1 \rangle + \frac{1}{16} \sum_{j=1}^N \frac{\phi_{2j}^2}{\lambda - \lambda_j}.$$

With respect to the standard Poisson bracket it is found that

$$\begin{aligned} \{A(\lambda), A(\mu)\} &= \{B(\lambda), B(\mu)\} = 0, \quad \{C(\lambda), C(\mu)\} = \frac{A(\lambda) - A(\mu)}{4}, \\ \{A(\lambda), B(\mu)\} &= \frac{B(\lambda) - B(\mu)}{8(\lambda - \mu)}, \quad \{A(\lambda), C(\mu)\} = \frac{C(\lambda) - C(\mu)}{8(\mu - \lambda)} - \frac{B(\lambda)}{8}, \\ \{B(\lambda), C(\mu)\} &= \frac{A(\lambda) - A(\mu)}{4(\lambda - \mu)}. \end{aligned} \quad (21)$$

It follows from (21) that

$$\{A(\lambda)^2 + B(\lambda)C(\lambda), A(\mu)^2 + B(\mu)C(\mu)\} = 0. \quad (22)$$

When $n = 3$, (17) for $\alpha = \frac{1}{32}$ yields the second higher-order constrained flow [23]

$$u_{xxxx} + 5u_x^2 + 10uu_{xx} + 10u^3 = \frac{1}{2} \langle \Phi_1, \Phi_1 \rangle, \quad (23a)$$

$$\phi_{1j,x} = \phi_{2j}, \quad \phi_{2j,x} = -(\lambda_j + u)\phi_{1j}, \quad j = 1, 2, \dots, N. \quad (23b)$$

Let $q_1 = u$, $q_2 = u_x$, $p_1 = -u_{xxx} - 10uu_x$, $p_2 = u_{xx}$, (23) becomes a FDIHS (19) with

$$H = \frac{1}{2}\langle\Phi_2, \Phi_2\rangle + \frac{1}{2}\langle\Lambda\Phi_1, \Phi_1\rangle + \frac{1}{2}q_1\langle\Phi_1, \Phi_1\rangle + \frac{1}{2}p_2^2 + q_2p_1 + 5q_1q_2^2 - \frac{5}{2}q_1^4.$$

We now consider the Rosochatius deformation $\tilde{N}^{(2)}$ of the Lax matrix $N^{(2)}$ with

$$\tilde{A}(\lambda) = A(\lambda), \quad \tilde{B}(\lambda) = B(\lambda), \quad \tilde{C}(\lambda) = C(\lambda) + \frac{1}{16} \sum_{j=1}^N \frac{\mu_j}{(\lambda - \lambda_j)\phi_{1j}^2}.$$

It is not difficult to find that $\tilde{A}(\lambda)$, $\tilde{B}(\lambda)$ and $\tilde{C}(\lambda)$ keep the relations of the Poisson brackets (21) and (22).

A direct calculation gives

$$\tilde{A}^2(\lambda) + \tilde{B}(\lambda)\tilde{C}(\lambda) = -\lambda^3 + P_0 + \sum_{j=1}^N \frac{P_j}{\lambda - \lambda_j} - \frac{1}{256} \sum_{j=1}^N \frac{\mu_j}{(\lambda - \lambda_j)^2}, \quad (24)$$

where

$$\begin{aligned} P_0 &= \frac{1}{16}(\langle\Phi_2, \Phi_2\rangle + \langle\Lambda\Phi_1, \Phi_1\rangle + q_1\langle\Phi_1, \Phi_1\rangle + 2q_1^3 + p_1^2 + \sum_{j=1}^N \frac{\mu_j}{\phi_{1j}^2}) \\ P_j &= \frac{p_1}{32}\phi_{1j}\phi_{2j} + \frac{1}{16}(\lambda_j - \frac{q_1}{2})(\phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2}) + \frac{1}{16}(\lambda_j^2 + \frac{q_1}{2}\lambda_j + \frac{1}{16}\langle\Phi_1, \Phi_1\rangle + \frac{q_1^2}{4})\phi_{1j}^2 \\ &+ \frac{1}{256} \sum_{k \neq j} \frac{1}{\lambda_j - \lambda_k} [2\phi_{1j}\phi_{1k}\phi_{2j}\phi_{2k} - \phi_{1j}^2(\phi_{2k}^2 + \frac{\mu_k}{\phi_{1k}^2}) - \phi_{1k}^2(\phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2})], \quad j = 1, \dots, N. \end{aligned} \quad (25)$$

Choosing $8P_0 = \tilde{H}$ as a Hamiltonian function, we get the following Hamiltonian system

$$q_{1x} = p_1, \quad p_{1x} = -\frac{1}{2}\langle\Phi_1, \Phi_1\rangle - 3q_1^2, \quad (26a)$$

$$\phi_{1jx} = \phi_{2j}, \quad \phi_{2jx} = -\lambda_j\phi_{1j} - q_1\phi_{1j} + \frac{\mu_j}{\phi_{1j}^3}, \quad (26b)$$

which is the Rosochatius deformation of the first higher-order constrained flow (20).

From (4), we have

$$\frac{d}{dx} \text{tr}(\tilde{N}^{(2)}(\lambda))^2 = \frac{d}{dx} [\tilde{A}^2(\lambda) + \tilde{B}(\lambda)\tilde{C}(\lambda)] = \text{tr}[U, (\tilde{N}^{(2)}(\lambda))^2] = 0, \quad (27)$$

which implies that $P_0, P_1, \dots, P_N$ are $N+1$ independent first integrals of the Hamiltonian system (26). The equality (22) of Poisson bracket for $\tilde{A}(\lambda)$, $\tilde{B}(\lambda)$ and $\tilde{C}(\lambda)$ indicates that $\{P_i, P_j\} = 0$, $i, j = 0, 1, \dots, N$. So the Rosochatius deformation (26) of the first higher-order constrained flow (20) is a FDIHS in the Liouville's sense [41].

Remark 1. For $N = 1$, $\lambda_1 = 0$, (26) yields

$$q_{1xx} = -\frac{1}{2}\phi_1^2 - 3q_1^2,$$

$$\phi_{1xx} = -q_1\phi_1 + \frac{\mu_1}{\phi_1^3},$$

which is the well-known generalized integrable Hénon-Heiles system [42]- [44]. In fact (26) can be regarded as the integrable multidimensional extension of Hénon-Heiles system.

Similarly, choosing

$$\tilde{H} = \frac{1}{2}\langle\Phi_2, \Phi_2\rangle + \frac{1}{2}\langle\Lambda\Phi_1, \Phi_1\rangle + \frac{1}{2}q_1\langle\Phi_1, \Phi_1\rangle + \frac{1}{2}p_2^2 + q_2p_1 + 5q_1q_2^2 - \frac{5}{2}q_1^4 + \frac{1}{2}\sum_{j=1}^N \frac{\mu_j}{\phi_{1j}^2},$$

we get the Rosochatius deformation of the second higher-order constrained flow (23)

$$q_{1x} = q_2, \quad q_{2x} = p_2, \quad p_{1x} = 10q_1^3 - 5q_2^2 - \frac{1}{2}\langle\Phi_1, \Phi_1\rangle, \quad p_{2x} = -10q_1q_2 - p_1, \quad (28a)$$

$$\phi_{1jx} = \phi_{2j}, \quad \phi_{2jx} = -\lambda_j\phi_{1j} - q_1\phi_{1j} + \frac{\mu_j}{\phi_{1j}^3}, \quad (28b)$$

which, in the same way, can be shown to be a FDIHS.

In general, the integrable Rosochatius deformation of the higher-order constrained flow (19) is generated by the following Hamiltonian function

$$\tilde{H} = \sum_{i=1}^{n-1} q_{i,x}p_i - H_n + \frac{1}{2}\langle\Phi_2, \Phi_2\rangle + \frac{1}{2}\langle\Lambda\Phi_1, \Phi_1\rangle + \frac{1}{2}q_1\langle\Phi_1, \Phi_1\rangle + \frac{1}{2}\sum_{j=1}^N \frac{\mu_j}{\phi_{1j}^2}. \quad (29)$$

The KdV hierarchy with self-consistent sources is defined by [25, 26, 29], [37]- [39]

$$u_{t_n} = \partial\left[\frac{\delta H_n}{\delta u} - \alpha \sum_{j=1}^N \frac{\delta \lambda_j}{\delta u}\right] \equiv \partial[-2b_{n+1} - \alpha \sum_{j=1}^N \phi_{1j}^2], \quad (30a)$$

$$\phi_{1jx} = \phi_{2j}, \quad \phi_{2j,x} = -(\lambda_j + u)\phi_{1j}, \quad j = 1, 2, \dots, N. \quad (30b)$$

Since the higher-order constrained flows (17) are just the stationary equation of the KdV hierarchy with self-consistent sources (30), it is obviously that the zero-curvature representation for the KdV hierarchy with self-consistent sources (30) is given by (6) with [29]

$$\begin{aligned} N^{(n)} = & \sum_{i=1}^n \begin{pmatrix} a_i & b_i \\ c_i & -a_i \end{pmatrix} \lambda^{n-i} + \begin{pmatrix} 0 & 0 \\ b_{n+1} + \frac{\alpha}{2} \sum_{j=1}^N \phi_{1j}^2 & 0 \end{pmatrix} \\ & + \frac{\alpha}{2} \sum_{j=1}^N \frac{1}{\lambda - \lambda_j} \begin{pmatrix} \phi_{1j}\phi_{2j} & -\phi_{1j}^2 \\ \phi_{2j}^2 & -\phi_{1j}\phi_{2j} \end{pmatrix}. \end{aligned} \quad (31)$$

Eq.(30) for $n = 2$, $\alpha = \frac{1}{8}$ gives rise to the KdV equation with self-consistent sources [29, 37]

$$u_t = -\frac{1}{4}(u_{xxx} + 6uu_x) - \frac{1}{8} \sum_{j=1}^N (\phi_{1j}^2)_x, \quad (32a)$$

$$\phi_{1jx} = \phi_{2j}, \quad \phi_{2jx} = -(\lambda_j + u)\phi_{1j}, \quad j = 1, 2, \dots, N. \quad (32b)$$

Based on (26), the Rosochatius deformation of KdV equation with self-consistent sources is given by

$$u_t = -\frac{1}{4}(u_{xxx} + 6uu_x) - \frac{1}{8} \sum_{j=1}^N (\phi_{1j}^2)_x, \quad (33a)$$

$$\phi_{1jx} = \phi_{2j}, \quad \phi_{2jx} = -(\lambda_j + u)\phi_{1j} + \frac{\mu_j}{\phi_{1j}^3}, \quad j = 1, \dots, N \quad (33b)$$

which has the zero-curvature representation (6) with the $N^{(2)}$ given by

$$N^{(2)} = \begin{pmatrix} \frac{u_x}{4} & \lambda - \frac{u}{2} \\ -\lambda^2 - \frac{u}{2}\lambda + \frac{1}{4}u_{xx} + \frac{1}{2}u^2 + \frac{1}{16} \sum_{j=1}^N \phi_{1j}^2 & -\frac{u_x}{4} \end{pmatrix} + \frac{1}{16} \sum_{j=1}^N \frac{1}{\lambda - \lambda_j} \begin{pmatrix} \phi_{1j}\phi_{2j} & -\phi_{1j}^2 \\ \phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2} & -\phi_{1j}\phi_{2j} \end{pmatrix}. \quad (34)$$

Remark 2: The fact that the stationary equation of (33) is a FDIHS (26) and (33) has zero-curvature representation (6) implies the integrability of the Rosochatius deformation of the KdV equation with self-consistent source (33).

In general the Rosochatius deformation of the KdV hierarchy with self-consistent sources is given by

$$u_{t_n} = \partial \left[\frac{\delta H_n}{\delta u} - \frac{2}{4^n} \sum_{j=1}^N \phi_{1j}^2 \right], \quad (35a)$$

$$\phi_{1jx} = \phi_{2j}, \quad \phi_{2j,x} = -(\lambda_j + u)\phi_{1j} + \frac{\mu_j}{\phi_{1j}^3}, \quad j = 1, 2, \dots, N. \quad (35b)$$

which has the zero-curvature representation (6) with $N^{(n)}$ given by

$$N^{(n)} = \sum_{i=1}^n \begin{pmatrix} a_i & b_i \\ c_i & -a_i \end{pmatrix} \lambda^{n-i} + \begin{pmatrix} 0 & 0 \\ b_{n+1} + \frac{1}{4^n} \sum_{j=1}^N \phi_{1j}^2 & 0 \end{pmatrix} + \frac{1}{4^n} \sum_{j=1}^N \frac{1}{\lambda - \lambda_j} \begin{pmatrix} \phi_{1j}\phi_{2j} & -\phi_{1j}^2 \\ \phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2} & -\phi_{1j}\phi_{2j} \end{pmatrix}.$$

3 The Rosochatius deformed AKNS hierarchy with self-consistent sources

For AKNS eigenvalue problem [40]

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}_x = U \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}, \quad U = \begin{pmatrix} -\lambda & q \\ r & \lambda \end{pmatrix},$$

and evolution equation of eigenfunction

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}_{t_n} = V^{(n)} \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}, \quad V^{(n)} = \sum_{i=1}^n \begin{pmatrix} a_i & b_i \\ c_i & -a_i \end{pmatrix} \lambda^{n-i},$$

the associated AKNS hierarchy reads

$$u_{t_n} = \begin{pmatrix} q \\ r \end{pmatrix}_{t_n} = J \begin{pmatrix} c_{n+1} \\ b_{n+1} \end{pmatrix} = J \frac{\delta H_{n+1}}{\delta u}$$

where

$$\begin{aligned} a_0 &= -1, \quad b_0 = c_0 = 0, \quad a_1 = 0, \quad b_1 = q, \quad c_1 = r, \dots, \\ \begin{pmatrix} c_{n+1} \\ b_{n+1} \end{pmatrix} &= L^n \begin{pmatrix} r \\ q \end{pmatrix}, \quad L = \frac{1}{2} \begin{pmatrix} \partial - 2r\partial^{-1}q & 2r\partial^{-1}r \\ -2q\partial^{-1}q & -\partial + 2q\partial^{-1}r \end{pmatrix}, \\ a_{n,x} &= qc_n - rb_n, \quad H_n = \frac{2}{n}a_{n+1}, \quad J = \begin{pmatrix} 0 & -2 \\ 2 & 0 \end{pmatrix}, \end{aligned}$$

we have

$$\frac{\delta \lambda}{\delta q} = \frac{1}{2}\phi_2^2, \quad \frac{\delta \lambda}{\delta r} = -\frac{1}{2}\phi_1^2.$$

The higher-order constrained flows of the AKNS hierarchy is given by [26,27]

$$\frac{\delta H_{n+1}}{\delta u} - \frac{1}{2} \sum_{j=1}^N \frac{\delta \lambda_j}{\delta u} = \begin{pmatrix} c_{n+1} \\ b_{n+1} \end{pmatrix} - \frac{1}{4} \begin{pmatrix} \langle \Phi_2, \Phi_2 \rangle \\ -\langle \Phi_1, \Phi_1 \rangle \end{pmatrix} = 0, \quad (36a)$$

$$\phi_{1jx} = -\lambda_j \phi_{1j} + q\phi_{2j}, \quad \phi_{2jx} = \lambda_j \phi_{2j} + r\phi_{1j}, \quad j = 1, 2, \dots, N. \quad (36b)$$

(36) for $n = 2$ gives the first higher-order constrained flow

$$-q_{xx} + 2q^2r - \sum_{j=1}^N \phi_{1j}^2 = 0, \quad r_{xx} - 2qr^2 - \sum_{j=1}^N \phi_{2j}^2 = 0, \quad (37a)$$

$$\phi_{1jx} = -\lambda_j \phi_{1j} + q\phi_{2j}, \quad \phi_{2jx} = \lambda_j \phi_{2j} + r\phi_{1j}, \quad j = 1, 2, \dots, N. \quad (37b)$$

Let $q_1 = q$, $q_2 = r$, $p_1 = -\frac{1}{2}r_x$, $p_2 = -\frac{1}{2}q_x$, $Q = (\phi_{11}, \phi_{12}, \dots, \phi_{1N}, q_1, q_2)^T$, $P = (\phi_{21}, \phi_{22}, \dots, \phi_{2N}, p_1, p_2)^T$. (37) becomes a FDIHS (19) with

$$H = -\langle \Lambda \Phi_1, \Phi_2 \rangle + \frac{q_1}{2} \langle \Phi_2, \Phi_2 \rangle - \frac{q_2}{2} \langle \Phi_1, \Phi_1 \rangle + \frac{1}{2} q_1^2 q_2^2 - 2p_1 p_2,$$

and has the Lax representation (4) with the entries of Lax matrix $N^{(2)}$ given by

$$A(\lambda) = -2\lambda^2 + q_1 q_2 + \frac{1}{2} \sum_{j=1}^N \frac{\phi_{1j} \phi_{2j}}{\lambda - \lambda_j}, \quad B(\lambda) = 2\lambda q_1 + 2p_2 - \frac{1}{2} \sum_{j=1}^N \frac{\phi_{1j}^2}{\lambda - \lambda_j},$$

$$C(\lambda) = 2\lambda q_2 - 2p_1 + \frac{1}{2} \sum_{j=1}^N \frac{\phi_{2j}^2}{\lambda - \lambda_j}.$$

With respect to the standard Poisson bracket it is found that

$$\begin{aligned}\{A(\lambda), A(\mu)\} &= \{B(\lambda), B(\mu)\} = \{C(\lambda), C(\mu)\} = 0, \quad \{A(\lambda), B(\mu)\} = \frac{B(\lambda) - B(\mu)}{\lambda - \mu}, \\ \{A(\lambda), C(\mu)\} &= \frac{C(\mu) - C(\lambda)}{\lambda - \mu}, \quad \{B(\lambda), C(\mu)\} = \frac{2[A(\lambda) - A(\mu)]}{\lambda - \mu},\end{aligned}\quad (38)$$

which gives rise to (22).

Now we consider the Rosochatius deformation $\tilde{N}^{(2)}$ of the Lax matrix $N^{(2)}$

$$\tilde{A}(\lambda) = A(\lambda), \quad \tilde{B}(\lambda) = B(\lambda), \quad \tilde{C}(\lambda) = C(\lambda) + \frac{1}{2} \sum_{j=1}^N \frac{\mu_j}{(\lambda - \lambda_j)\phi_{1j}^2}. \quad (39)$$

It is easy to find that the elements in $\tilde{N}^{(2)}$ still keep the relations of the Poisson brackets (38) and (22).

A direct calculation gives

$$\tilde{A}^2(\lambda) + \tilde{B}(\lambda)\tilde{C}(\lambda) = 4\lambda^4 + P_0\lambda + P_1 + \sum_{j=1}^N \frac{P_j}{\lambda - \lambda_j} - \frac{1}{4} \sum_{j=1}^N \frac{\mu_j}{(\lambda - \lambda_j)^2} \quad (40)$$

where

$$\begin{aligned}P_0 &= -4q_1p_1 + 4q_2p_2 - 2\langle\Phi_1, \Phi_2\rangle, \\ P_1 &= -4p_1p_2 + q_1^2q_2^2 + q_1(\langle\Phi_2, \Phi_2\rangle + \sum_{j=1}^N \frac{\mu_j}{\phi_{1j}^2}) - q_2\langle\Phi_1, \Phi_1\rangle - 2\langle\Lambda\Phi_1, \Phi_2\rangle, \\ P_{j+1} &= (\lambda_jq_1 + p_2)(\phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2}) - (\lambda_jq_2 - p_1)\phi_{1j}^2 + (q_1q_2 - 2\lambda_j^2)\phi_{1j}\phi_{2j} \\ &+ \frac{1}{4} \sum_{k \neq j} \frac{1}{\lambda_j - \lambda_k} [2\phi_{1j}\phi_{2j}\phi_{1k}\phi_{2k} - \phi_{1j}^2(\phi_{2k}^2 + \frac{\mu_k}{\phi_{1k}^2}) - \phi_{1k}^2(\phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2})], \quad j = 1, \dots, N.\end{aligned}\quad (41)$$

Choosing $\frac{1}{2}P_1 = \tilde{H}$ as a Hamiltonian function, we get the following Hamiltonian system

$$q_{1x} = -2p_2, \quad q_{2x} = -2p_1, \quad (42a)$$

$$p_{1x} = -\frac{1}{2}\langle\Phi_2, \Phi_2\rangle - \sum_{j=1}^N \frac{\mu_j}{2\phi_{1j}^3} - q_2^2q_1, \quad p_{2x} = \frac{1}{2}\langle\Phi_1, \Phi_1\rangle - q_1^2q_2, \quad (42b)$$

$$\phi_{1jx} = -\lambda_j\phi_{1j} + q_1\phi_{2j}, \quad \phi_{2jx} = \lambda_j\phi_{2j} + q_2\phi_{1j} + \frac{\mu_jq_1}{\phi_{1j}^3}, \quad j = 1, \dots, N \quad (42c)$$

which has the Lax representation (4) with $N^{(2)}$ replaced by $\tilde{N}^{(2)}$. Then (22) and (27) implies that $P_0, P_1, \dots, P_{N+1}$ are $N+2$ independent first integrals in involution, so (42) is a FDIHS [41].

In general, in the similar way as in Sec.2, (36) becomes a FDIHS (19) with

$$H = \sum_{i=1}^n q_{i,x} p_i - H_{n+1} - \langle \Lambda \Phi_1, \Phi_2 \rangle + \frac{q_1}{2} \langle \Phi_2, \Phi_2 \rangle - \frac{q_2}{2} \langle \Phi_1, \Phi_1 \rangle.$$

Then the integrable Rosochatius deformation of the higher-order constrained flow (36) is generated by the following Hamiltonian function

$$\tilde{H} = \sum_{i=1}^n q_{i,x} p_i - H_{n+1} - \langle \Lambda \Phi_1, \Phi_2 \rangle + \frac{q_1}{2} \langle \Phi_2, \Phi_2 \rangle - \frac{q_2}{2} \langle \Phi_1, \Phi_1 \rangle + \frac{1}{2} \sum_{j=1}^N \frac{\mu_j q_1}{\phi_{1j}^2}. \quad (43)$$

The AKNS hierarchy with self-consistent sources is [26]

$$\begin{pmatrix} q \\ r \end{pmatrix}_{t_n} = J \left[\frac{\delta H_{n+1}}{\delta u} - \frac{1}{2} \sum_{j=1}^N \frac{\delta \lambda_j}{\delta u} \right] = J \left[\begin{pmatrix} c_{n+1} \\ b_{n+1} \end{pmatrix} - \frac{1}{4} \begin{pmatrix} \langle \Phi_2, \Phi_2 \rangle \\ -\langle \Phi_1, \Phi_1 \rangle \end{pmatrix} \right], \quad (44a)$$

$$\phi_{1jx} = -\lambda_j \phi_{1j} + q \phi_{2j}, \quad \phi_{2jx} = \lambda_j \phi_{2j} + r \phi_{1j}, \quad j = 1, 2, \dots, N. \quad (44b)$$

When $n = 2$, the AKNS equation with self-consistent sources reads [26]

$$2q_t = -q_{xx} + 2q^2 r - \sum_{j=1}^N \phi_{1j}^2, \quad 2r_t = r_{xx} - 2qr^2 - \sum_{j=1}^N \phi_{2j}^2, \quad (45a)$$

$$\phi_{1jx} = -\lambda_j \phi_{1j} + q \phi_{2j}, \quad \phi_{2jx} = \lambda_j \phi_{2j} + r \phi_{1j}, \quad j = 1, \dots, N. \quad (45b)$$

Based on (42), we obtain the integrable Rosochatius deformation of the AKNS equation with self-consistent sources

$$2q_t = -q_{xx} + 2q^2 r - \sum_{j=1}^N \phi_{1j}^2, \quad 2r_t = r_{xx} - 2qr^2 - \sum_{j=1}^N \phi_{2j}^2 - \sum_{j=1}^N \frac{\mu_j}{\phi_{1j}^2}, \quad (46a)$$

$$\phi_{1jx} = -\lambda_j \phi_{1j} + q \phi_{2j}, \quad \phi_{2jx} = \lambda_j \phi_{2j} + r \phi_{1j} + \frac{\mu_j q}{\phi_{1j}^3}, \quad j = 1, \dots, N \quad (46b)$$

which has the zero-curvature representation (6) with $N^{(2)}$

$$N^{(2)} = \begin{pmatrix} -2\lambda^2 + qr & 2\lambda q - q_x \\ 2\lambda r + r_x & 2\lambda^2 - qr \end{pmatrix} + \frac{1}{2} \sum_{j=1}^N \frac{1}{\lambda - \lambda_j} \begin{pmatrix} \phi_{1j} \phi_{2j} & -\phi_{1j}^2 \\ \phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2} & -\phi_{1j} \phi_{2j} \end{pmatrix}. \quad (47)$$

In general the integrable Rosochatius deformation of the AKNS hierarchy with self-consistent sources is given by

$$\begin{pmatrix} q \\ r \end{pmatrix}_{t_n} = J \left[\begin{pmatrix} c_{n+1} \\ b_{n+1} \end{pmatrix} - \frac{1}{4} \begin{pmatrix} \langle \Phi_2, \Phi_2 \rangle + \sum_{j=1}^N \frac{\mu_j}{\phi_{1j}^2} \\ -\langle \Phi_1, \Phi_1 \rangle \end{pmatrix} \right], \quad (48a)$$

$$\phi_{1jx} = -\lambda_j \phi_{1j} + q \phi_{2j}, \quad \phi_{2jx} = \lambda_j \phi_{2j} + r \phi_{1j} + \frac{\mu_j q}{\phi_{1j}^3}, \quad j = 1, 2, \dots, N. \quad (48b)$$

which has the zero-curvature representation (6) with $N^{(n)}$ given by

$$N^{(n)} = V^{(n)} + \frac{1}{2} \sum_{j=1}^N \frac{1}{\lambda - \lambda_j} \begin{pmatrix} \phi_{1j}\phi_{2j} & -\phi_{1j}^2 \\ \phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2} & -\phi_{1j}\phi_{2j} \end{pmatrix}.$$

Remark 3. *In contrast with the Rosochatius deformation of KdV hierarchy with self-consistent sources, the Rosochatius deformation of AKNS hierarchy with self-consistent sources has the deformed term in both (48a) and (48b).*

4 The Rosochatius deformed mKdV hierarchy with self-consistent sources

For mKdV spectral problem [32]

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}_x = U \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}, \quad U = \begin{pmatrix} -u & \lambda \\ \lambda & u \end{pmatrix},$$

and evolution equation of eigenfunction

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}_{t_n} = V^{(n)} \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}, \quad V^{(n)} = \sum_{i=1}^{n-1} \begin{pmatrix} a_i \lambda & b_i \\ c_i & -a_i \lambda \end{pmatrix} \lambda^{2n-2i-3} + \begin{pmatrix} a_n & 0 \\ 0 & -a_n \end{pmatrix},$$

the associated mKdV hierarchy reads

$$u_{t_n} = -\partial a_n = \partial \frac{\delta H_n}{\delta u},$$

where

$$a_0 = 0, \quad b_0 = c_0 = 1, \quad a_1 = -u, \quad b_1 = -\frac{u^2}{2} + \frac{u_x}{2}, \quad c_1 = -\frac{u^2}{2} - \frac{u_x}{2}, \dots,$$

$$a_{n+1} = L a_n, \quad L = \frac{1}{4} \partial^2 - u \partial^{-1} u \partial$$

$$b_n = \partial^{-1} u \partial a_n - \frac{1}{2} a_{nx}, \quad c_n = \partial^{-1} u \partial a_n + \frac{1}{2} a_{nx},$$

we have

$$\frac{\delta \lambda}{\delta u} = \frac{1}{2} \phi_1 \phi_2.$$

The higher-order constrained flows of the mKdV hierarchy is

$$\frac{\delta H_n}{\delta u} + 2 \sum_{j=1}^N \frac{\delta \lambda_j}{\delta u} \equiv -a_n + \sum_{j=1}^N \phi_{1j} \phi_{2j} = 0, \quad (49a)$$

$$\phi_{1j,x} = -u \phi_{1j} + \lambda_j \phi_{2j}, \quad \phi_{2j,x} = \lambda_j \phi_{1j} + u \phi_{2j}, \quad j = 1, 2, \dots, N. \quad (49b)$$

When $n = 2$, (49) gives the first higher-order constrained flow

$$u_{xx} - 2u^3 = -4 \sum_{j=1}^N \phi_{1j} \phi_{2j} = -4 \langle \Phi_1, \Phi_2 \rangle, \quad (50a)$$

$$\phi_{1jx} = \lambda_j \phi_{2j} - u \phi_{1j}, \quad \phi_{2jx} = \lambda_j \phi_{1j} + u \phi_{2j}, \quad j = 1, \dots, N. \quad (50b)$$

Let $q_1 = u$, $p_1 = -\frac{u_x}{4}$, it becomes a FDIHS (19) with

$$H = -q_1 \langle \Phi_1, \Phi_2 \rangle + \frac{1}{2} \langle \Lambda \Phi_2, \Phi_2 \rangle - \frac{1}{2} \langle \Lambda \Phi_1, \Phi_1 \rangle - 2p_1^2 + \frac{1}{8} q_1^4,$$

and has the Lax representation (4) with the entries of Lax matrix $N^{(2)}$ given by

$$A(\lambda) = -q_1 \lambda + \sum_{j=1}^N \frac{\lambda \phi_{1j} \phi_{2j}}{\lambda^2 - \lambda_j^2}, \quad B(\lambda) = \lambda^2 - \frac{1}{2} q_1^2 - 2p_1 - \sum_{j=1}^N \frac{\lambda_j \phi_{1j}^2}{\lambda^2 - \lambda_j^2}, \quad (51)$$

$$C(\lambda) = \lambda^2 - \frac{1}{2} q_1^2 + 2p_1 + \sum_{j=1}^N \frac{\lambda_j \phi_{2j}^2}{\lambda^2 - \lambda_j^2}.$$

With respect to the standard Poisson bracket, a direct calculation gives

$$\begin{aligned} \{A(\lambda), A(\mu)\} &= \{B(\lambda), B(\mu)\} = \{C(\lambda), C(\mu)\} = 0, \quad \{A(\lambda), B(\mu)\} = 2\lambda \frac{B(\lambda) - B(\mu)}{\lambda^2 - \mu^2}, \\ \{A(\lambda), C(\mu)\} &= 2\lambda \frac{C(\lambda) - C(\mu)}{\mu^2 - \lambda^2}, \quad \{B(\lambda), C(\mu)\} = \frac{4[A(\mu)\mu - A(\lambda)\lambda]}{\mu^2 - \lambda^2} \end{aligned} \quad (52)$$

which leads to (22).

Now we consider the integrable Rosochatius deformation $\tilde{N}^{(2)}$ of the Lax matrix $N^{(2)}$

$$\tilde{A}(\lambda) = A(\lambda), \quad \tilde{B}(\lambda) = B(\lambda), \quad \tilde{C}(\lambda) = C(\lambda) + \sum_{j=1}^N \frac{\mu_j \lambda_j}{(\lambda^2 - \lambda_j^2) \phi_{1j}^2}. \quad (53)$$

It is not difficult to find that the elements in $\tilde{N}^{(2)}$ still keep the relations of the Poisson brackets (52) and (22).

A direct calculation gives

$$\tilde{A}^2(\lambda) + \tilde{B}(\lambda) \tilde{C}(\lambda) = \lambda^4 + P_0 + \sum_{j=1}^N \frac{P_j}{\lambda^2 - \lambda_j^2} - \sum_{j=1}^N \frac{\lambda_j \mu_j}{(\lambda^2 - \lambda_j^2)^2} \quad (54)$$

where

$$\begin{aligned} P_0 &= -2q_1 \langle \Phi_1, \Phi_2 \rangle + \langle \Lambda \Phi_2, \Phi_2 \rangle - \langle \Lambda \Phi_1, \Phi_1 \rangle + \sum_{j=1}^N \frac{\lambda_j \mu_j}{\phi_{1j}^2} + \frac{1}{4} q_1^4 - 4p_1^2 \\ P_j &= -2q_1 \lambda_j^2 \phi_{1j} \phi_{2j} + \lambda_j^3 (\phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2}) - (\frac{1}{2} q_1^2 + 2p_1) (\lambda_j \phi_{2j}^2 + \frac{\lambda_j \mu_j}{\phi_{1j}^2}) - \lambda_j^3 \phi_{1j}^2 + (\frac{1}{2} q_1^2 - 2p_1) \lambda_j \phi_{1j}^2 \end{aligned} \quad (55)$$

$$-\sum_{k \neq j} \frac{1}{\lambda_j^2 - \lambda_k^2} [2\lambda_j^2 \phi_{1j} \phi_{2j} \phi_{1k} \phi_{2k} + \lambda_j \lambda_k \phi_{1j}^2 (\phi_{2k}^2 + \frac{\mu_k}{\phi_{1k}^2}) + \lambda_j \lambda_k \phi_{1k}^2 (\phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2})].$$

Choosing $\frac{1}{2}P_0 = \tilde{H}$ as Hamiltonian function, we get the following Hamiltonian system

$$q_{1x} = -4p_1, \quad p_{1x} = \langle \Phi_1, \Phi_2 \rangle - \frac{1}{2}q_1^3, \quad (56a)$$

$$\phi_{1jx} = \lambda_j \phi_{2j} - q_1 \phi_{1j}, \quad \phi_{2jx} = \lambda_j \phi_{1j} + q_1 \phi_{2j} + \frac{\lambda_j \mu_j}{\phi_{1j}^3}, \quad j = 1, \dots, N \quad (56b)$$

which has the Lax representation (4) with $N^{(2)}$ replaced by $\tilde{N}^{(2)}$ and is a FDIHS.

Similarly, (49) can be transformed into a FDIHS (19) with $N + 1$ independent first integrals $P_0, P_1, \dots, P_N$ in involution

$$H = \sum_{i=1}^{n-1} q_{i,x} p_i - H_n - q_1 \langle \Phi_1, \Phi_2 \rangle + \frac{1}{2} \langle \Lambda \Phi_2, \Phi_2 \rangle - \frac{1}{2} \langle \Lambda \Phi_1, \Phi_1 \rangle.$$

Then the integrable Rosochatius deformation of the higher-order constrained flow (49) is generated by the following Hamiltonian function

$$\tilde{H} = \sum_{i=1}^{n-1} q_{i,x} p_i - H_n - q_1 \langle \Phi_1, \Phi_2 \rangle + \frac{1}{2} \langle \Lambda \Phi_2, \Phi_2 \rangle - \frac{1}{2} \langle \Lambda \Phi_1, \Phi_1 \rangle + \frac{1}{2} \sum_{j=1}^N \frac{\lambda_j \mu_j}{\phi_{1j}^2}. \quad (57)$$

The mKdV hierarchy with self-consistent sources is

$$u_{tn} = \partial \left[\frac{\delta H_n}{\delta u} + 2 \sum_{j=1}^N \frac{\delta \lambda_j}{\delta u} \right] \equiv \partial [-a_n + \sum_{j=1}^N \phi_{1j} \phi_{2j}], \quad (58a)$$

$$\phi_{1j,x} = -u \phi_{1j} + \lambda_j \phi_{2j}, \quad \phi_{2j,x} = \lambda_j \phi_{1j} + u \phi_{2j}, \quad j = 1, 2, \dots, N. \quad (58b)$$

When $n = 2$, (58) gives the mKdV equation with self-consistent sources

$$u_t = \frac{u_{xxx}}{4} - \frac{3}{2} u^2 u_x + \sum_{j=1}^N (\phi_{1j} \phi_{2j})_x, \quad (59a)$$

$$\phi_{1jx} = \lambda_j \phi_{2j} - u \phi_{1j}, \quad \phi_{2jx} = \lambda_j \phi_{1j} + u \phi_{2j}, \quad j = 1, \dots, N. \quad (59b)$$

Based on (56), the integrable Rosochatius deformation of the mKdV equation with self-consistent sources is given by

$$u_t = \frac{u_{xx}}{4} - \frac{3}{2} u^2 u_x + \sum_{j=1}^N (\phi_{1j} \phi_{2j})_x, \quad (60a)$$

$$\phi_{1jx} = \lambda_j \phi_{2j} - u \phi_{1j}, \quad \phi_{2jx} = \lambda_j \phi_{1j} + u \phi_{2j} + \frac{\lambda_j \mu_j}{\phi_{1j}^3}, \quad j = 1, \dots, N. \quad (60b)$$

which has the zero-curvature representation (6) with $N^{(2)}$ given by

$$N^{(2)} = \begin{pmatrix} -u\lambda^2 - \frac{u_{xx}}{4} + \frac{u^3}{2} & \lambda^3 - (\frac{u^2}{2} - \frac{u_x}{2})\lambda \\ \lambda^3 - (\frac{u^2}{2} + \frac{u_x}{2})\lambda & u\lambda^2 + \frac{u_{xx}}{4} - \frac{u^3}{2} \end{pmatrix} + \begin{pmatrix} -\sum_{j=1}^N \phi_{1j}\phi_{2j} & 0 \\ 0 & \sum_{j=1}^N \phi_{1j}\phi_{2j} \end{pmatrix} \\ + \sum_{j=1}^N \frac{\lambda}{\lambda^2 - \lambda_j^2} \begin{pmatrix} \lambda\phi_{1j}\phi_{2j} & -\lambda_j\phi_{1j}^2 \\ \lambda_j(\phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2}) & -\lambda\phi_{1j}\phi_{2j} \end{pmatrix} \quad (61)$$

In general the integrable Rosochatius deformation of the mKdV hierarchy with self-consistent sources is given by

$$u_{t_n} = \partial \left[\frac{\delta H_n}{\delta u} + 2 \sum_{j=1}^N \frac{\delta \lambda_j}{\delta u} \right] = \partial [-a_n + \sum_{j=1}^N \phi_{1j}\phi_{2j}] \quad (62a)$$

$$\phi_{1jx} = \lambda_j\phi_{2j} - u\phi_{1j}, \quad \phi_{2jx} = \lambda_j\phi_{1j} + u\phi_{2j} + \frac{\lambda_j\mu_j}{\phi_{1j}^3}, \quad j = 1, \dots, N. \quad (62b)$$

which has the zero-curvature representation (6) with $N^{(n)}$ given by

$$N^{(n)} = \sum_{i=1}^{n-1} \begin{pmatrix} a_i\lambda & b_i \\ c_i & -a_i\lambda \end{pmatrix} \lambda^{2n-2i-3} + \begin{pmatrix} a_n - \sum_{j=1}^N \phi_{1j}\phi_{2j} & 0 \\ 0 & a_n + \sum_{j=1}^N \phi_{1j}\phi_{2j} \end{pmatrix} \\ + \sum_{j=1}^N \frac{\lambda}{\lambda^2 - \lambda_j^2} \begin{pmatrix} \lambda\phi_{1j}\phi_{2j} & -\lambda_j\phi_{1j}^2 \\ \lambda_j(\phi_{2j}^2 + \frac{\mu_j}{\phi_{1j}^2}) & -\lambda\phi_{1j}\phi_{2j} \end{pmatrix}.$$

5 Conclusion

Rosochatius-type integrable systems have important physical applications. However, the studies on Rosochatius deformation are limited to few finite dimensional integrable Hamiltonian system(FDIHS) at present. The main purpose of this paper is to propose a systematic method to generalize the Rosochatius deformation for FDIHS to the Rosochatius deformation for infinite dimensional integrable equations. We first construct infinite number of integrable Rosochatius deformations of FDIHSs obtained from higher-order constrained flows of some soliton hierarchies. Then, based on the Rosochatius deformed higher-order constrained flows, we establish the integrable Rosochatius deformations of some soliton hierarchies with self-consistent sources. The Rosochatius deformations of the KdV hierarchy with self-consistent sources, of the AKNS hierarchy with self-consistent sources and of the mKdV hierarchy with self-consistent sources, together with their Lax pairs are obtained. The approach presented here can be applied to other cases.

Acknowledgments

This work is supported by National Basic Research Program of China (973 Program) (2007CB814800) and National Natural Science Foundation of China (10671121).

References

- [1] Rosochatius E, 1877 dissertation, University of Göttingen.
- [2] Neumann C, 1859 J. Reine Angew. Math. **56** 46.
- [3] Wojciechowski S, 1985 Phys. Scr. **31** 433.
- [4] Tondo G, 1995 J. Phys. A **28** 5097.
- [5] Kubo R, Ogura W, Saito T and Yasui Y, 1999 Phys. Lett. A **251** 6.
- [6] Jacobi C G, 1884 Vorlesungen über dynamik, (Verlag G. Reimer, Berlin).
- [7] Moser J, Geometry of Quadrics and Spectral Theory. The Chern Symposium 1979. Proc. Internat. Sympos. Berkeley, California, 1979 (Springer, New York-Berlin).
- [8] Knörrer H, 1982 J. Reine Angew. Math. **334** 69.
- [9] Bartocci C, Falqui G and Pedroni M, 2004 Diff. Geom. Applic. **21** 349 .
- [10] Adams M R, Harnad J and Hurtubise J, 1993 Commun. Math. Phys. **155** 385.
- [11] Arutyunov G, Russo J and Tseytlin A A, 2004 Phys. Rev. D **69** 086009.
- [12] Harnad J and Winternitz P, 1995 Commun. Math. Phys. **172** 263.
- [13] Arutyunov G, Russo J and Tseytlin A A, 2003 Nucl. Phys. B **671** 3.
- [14] Kruczenski M, Russo J G and Tseytlin A A, 2006 J. High Energy Phys. **10** 063.
- [15] McLoughlin T and Wu X K, 2006 J. High Energy Phys. **8** 063.
- [16] Dimov H and Rashkov R, 2005 Int. J. Mod. Phys. A **20** 4337.
- [17] Bozhilov P, 2007 J. High Energy Phys. **8** 073.
- [18] Christiansen P L, Eilbeck J C, Enolskii V Z and Kostov N A, 2000 Proc. R. Soc, London, Ser. A **456** 2263.
- [19] Zhou R G, 2007 J. Math. Phys. **48** 103510.
- [20] Cao C W, 1990 Sci China, Ser A: Math. Phys., Astron, Technol. Sci **33** 528.
- [21] Cao C W and Geng X G, 1991 J. Math. Phys. **32** 2223.

- [22] Zeng Y B and Li Y S, 1990 J. Phys. A: Math. Gen. **23** L89.
- [23] Zeng Y B, 1991 Phys. Lett. A **160** 541.
- [24] Zeng Y B and Li Y S, 1993 J. Phys. A: Math. Gen. **26** L273.
- [25] Zeng Y B , 1994 Physica D, **73** 171.
- [26] Zeng Y B, 1995 Acta Mathematica Scientia, **15** 337.
- [27] Zeng Y B and Li Y S, 1996 Acta Math. Appl. Sinica **12** 337.
- [28] Zeng Y B and Hietarinta J, 1996 J. Phys. A: Math. Gen. **29** 5241.
- [29] Zeng Y B, Ma W X and Shao Y J, 2001 J. Math. Phys. **42** 2113.
- [30] Zeng Y B, Shao Y J and Ma W X, 2002 Commun. Theor. Phys. **38** 641.
- [31] Antownowicz M and Wojciechowski S R, 1990 Phys. Lett. A **147** 455.
- [32] Zhou R G and Ma W X, 1998 J. Phys. Soc. Jpn. **67** 4045.
- [33] Ma W X and Zhou R G , 2001 J. Phys. A **34** 867.
- [34] Deleón M and Rodrigues P R, 1985 Generalized classical mechanics and field theory (North-Holland, Amsterdam).
- [35] Blaszkak M, J.Math.Phys. 1995 **36** 4826.
- [36] Zeng Y B, Physica A 1999 **262** 405.
- [37] Mel'nikov V K, 1989 Commun. Math. Phys. **120** 451.
- [38] Mel'nikov V K, 1990 Inverse Probl., **6** 233.
- [39] Mel'nikov V K, 1990 J. Math. Phys. **31** 1106.
- [40] Ablowitz M J and Segur H, 1991 Solitons and the Inverse Scattering Transformation, (SIAM, Philadelphia).
- [41] Arnold V I, 1978 Mathematical Method of Classical Mechanics (Springer-Verlag, New York).
- [42] Hénon M and Heiles C, 1964 Astron J. **69** 73.
- [43] Chang Y F, Tabor M and Weiss J, 1982 J. Math. Phys. **23** 531.
- [44] Antownowicz M and Wojciechowski S R, 1992 Phys. Lett. A **163** 167.